

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/12, 15/54, A01K 67/027, C12Q 1/48, 1/68, A61K 38/45	A1	(11) International Publication Number: WO 97/42310 (43) International Publication Date: 13 November 1997 (13.11.97)
(21) International Application Number: PCT/DK97/00200 (22) International Filing Date: 2 May 1997 (02.05.97) (30) Priority Data: 0539/96 6 May 1996 (06.05.96) DK (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK) (72) Inventors; and (75) Inventors/Applicants (for US only): HANSEN, Torben [DK/DK]; Helleruplund Allé 3, DK-2900 Hellerup (DK). ANDERSEN, Carsten, Bo [DK/US]; 907 Manor Way, Los Altos, CA 94024 (US). PEDERSEN, Oluf, Borbye [DK/DK]; Rydholt 11, DK-2840 Holte (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsvaerd (DK).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: MUTANT cDNA ENCODING THE p85 α SUBUNIT OF PHOSPHATIDYLINOSITOL 3-KINASE (57) Abstract A nucleic acid sequence comprising a DNA sequence encoding a regulatory subunit of phosphatidylinositol 3-kinase (P13K), the DNA sequence containing a mutation of at least one nucleotide, or comprising a segment of the DNA sequence including said mutation for use as a diagnostic tool, a marker or a probe. Examples of mutations in p85 α subunit are: C to T in nucleotide 261; T to G in nucleotide 663; A to G in nucleotide 810; G to A in nucleotide 1020.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MUTANT cDNA ENCODING THE p85 α SUBUNIT OF PHOSPHATIDYLINOSITOL 3-KINASE

5

FIELD OF THE INVENTION

The present invention relates to a mutant cDNA sequence encoding the regulatory p85 α subunit of phosphatidylinositol 3-kinase (PI3K), a method of detecting a mutation in the
10 gene encoding the regulatory p85 α subunit of phosphatidylinositol 3-kinase, as well as a diagnostic composition and a test kit for use in the method.

BACKGROUND OF THE INVENTION

15 Phosphatidylinositol 3-kinase is a heterodimeric protein consisting of a catalytic 110 kDa subunit and an α -85 kDa subunit that is necessary for docking to growth factor plasma membrane receptors including the insulin receptor and to signalling phosphotyrosine motifs in signalling proteins, e.g. insulin receptor substrate-1 (1,2). In addition to its association with receptors that mediate their effects via tyrosine kinase activity, PI 3-kinases are also
20 implicated in facilitating membrane trafficking processes. The pinocytic activity, membrane ruffling and actin reorganization response observed when KB cells are exposed to insulin have been shown to be due to PI3-kinase activation (3,4). PI3-kinase has been implicated as a key intermediate in the cytoskeletal rearrangements that accompany secretory processes in platelets, basophil and neutrophil cells (5,6). Similarly, the yeast homologue VPS34 has
25 been shown to be involved in vesicle trafficking and protein sorting (7). PI3-kinase is thus an attractive candidate protein as a mediator of insulin action on glucose transport as it could act as a point of convergence of signalling and trafficking processes.

In our attempts to elucidate the genetics of insulin action and glucose metabolism, we have
30 performed mutational analysis on the regulatory p85 α subunit of PI3K which is expressed in skeletal muscle.

References:

1. Hiles, I.D., Otsu, M., Volinia, S., Fry, M.J., Gout, I., Dhand, R., Panayotou, G., Ruiz-Larrea, F., Thompson, A., Totty, N.F., Hsuan, J.J., Courtneige, S.A., Parker, P.J., and Waterfield, M.D. (1992) *Cell* **70**, 419-429.
- 5 2. Myers, M.G., and White, M.F. (1993) *Diabetes* **42**, 643-650.
3. Kotani, K., Yonezawa, K., Hara, K., Ueda, H., Kitamura, Y., Sakaue, H., Ando, A., Chavanieu, A., Calas, b., Grigorescu, f., Nishiyama, M., Waterfield, m.d., and Kasuga, M. (1994) *EMBO J.* **13**, 2313-2321.
4. Kotani, K., Hara, K., Kotani, K., Yonezawa, K., and Kasuga, M. (19995) *Biochem.*
- 10 *Biophys. Res. Comm.* (in press).
5. Yatomi, Y., Hazeki, O., Kume, S., and Ui, M. (11111992) *Biochem. J.* **285**, 745-751.
6. Yano, H., Nakanishi, S., Kimura, K., Hanai, N., Saiitoh, Y., Fukui, Y., Nomomura, Y., and Matsuda, Y. (1993) *J. Biol. Chem.* **268**, 25846-25856.
7. Schu, P.V., Kaoru, T., Fry, M.J., Stack, J.H., Waterfield, M.D., and Emr, S.D.
- 15 (1993) *Science* **260**, 88-91.

SUMMARY OF THE INVENTION

The gene encoding the regulatory p85 α subunit of phosphatidylinositol 3-kinase has been
20 examined for genetic variations which might be associated with or causing impaired insulin
sensitivity, impaired glucose effectiveness, and/or impaired glucose disappearance.

The present invention relates to a nucleic acid sequence comprising a DNA sequence
encoding the regulatory p85 α subunit of human phosphatidylinositol 3-kinase (PI3K), the
25 DNA sequence containing a mutation of at least one nucleotide, or comprising a segment of
the cDNA sequence including said mutation. It has been found that such mutations may be
localized to the p85 α subunit of PI3K.

It has been found that mutation of the regulatory p85 α subunit of PI3K gene may be
30 indicative of abnormalities significant for the development of changes in whole body
glucose metabolism and whole body insulin action. For instance, the mutation may give rise
to the substitution of an amino acid in the regulatory p85 α subunit of PI3K which may

cause changes in the structure, such as the tertiary structure, or biochemical properties of the regulatory p85 α subunit of PI3K. Such changes may interfere with the normal binding of the regulatory p85 α subunit of PI3K to the insulin receptor kinase and insulin receptor substrates, e.g. IRS-1 and IRS-2 leading to altered glucose metabolism and reduced insulin sensitivity, and may lead to the development of insulin resistant conditions, such as NIDDM, cardiovascular diseases, obesity, and hypertension in human beings and other mammals.

In another aspect, the present invention relates to a living system containing a nucleic acid sequence of the invention and capable of expressing the regulatory p85 α subunit of PI3K wherein at least one amino acid is substituted, inserted or deleted, or the living system contains a nucleic acid sequence of the invention carrying at least one silent mutation. The living system, which may comprise a cell or a multicellular organism containing the appropriate signal transduction pathway, may be used to screen for substances which have an effect on the regulatory p85 α subunit of PI3K activity.

In a further aspect, the present invention relates to a method of detecting the presence of a mutation in the gene encoding the regulatory p85 α subunit of PI3K, the method comprising obtaining a biological sample from a subject and analysing the sample for a mutation of at least one nucleotide. It has now been shown that homozygous carriers of a the regulatory p85 α subunit of PI3K mutation G to A in nucleotide # 1020 exhibit a decreased glucose disappearance rate and a decreased glucose effectiveness and a decreased insulin sensitivity. Therefore, the present method is useful in diagnosing predisposition to insulin resistance and possibly impaired glucose tolerance and related diseases, such as NIDDM, cardiovascular diseases, obesity, and hypertension, in a subject as well as other disorders resulting from an altered glucose metabolism. Biological samples may, for instance, be obtained from blood, serum, plasma or tissue. The invention further relates to a diagnostic composition and a test kit for use in the method.

DETAILED DESCRIPTION OF THE INVENTION

Examples of specific mutations found in the cDNA sequence encoding the regulatory p85 α subunit of PI3K are:

C to T in nucleotide # 261;

T to G in nucleotide # 663;

A to G in nucleotide # 810;

G to A in nucleotide # 1020

10 according to the numbering of E.Y. Skolnik et al., *Cell* 65, 1991, pp. 83-90.

It is at present assumed that mutations giving rise to substitutions of one or more amino acids in the regulatory p85 α subunit of PI3K may be of particular importance as they may cause changes in the tertiary structure of the regulatory p85 α subunit of PI3K and thereby
15 change its biological activity. Consequently, in a particular embodiment, the present invention relates to a nucleic acid sequence comprising a cDNA sequence encoding the regulatory p85 α subunit of PI3K and containing a mutation giving rise to at least one amino acid substitution in the regulatory p85 α subunit of PI3K protein sequence. The mutation may for instance be located at a site where the amino acid substitution interferes with the
20 binding of the regulatory p85 α subunit of PI3K to insulin receptor kinase or insulin signalling proteins, e.g. IRS-1 and/or IRS-2, as such a mutation is most likely to be involved in changes in insulin signalling and basal and insulin stimulated glucose uptake. Such changes may therefore lead to reductions in whole body insulin sensitivity and whole body glucose metabolism. Martin, B.C. et al. (*Lancet* 1992, Vol. 340, 925:929) have
25 previously shown that individuals having low or reduced Si (insulin sensitivity) or Sg (glucose effectiveness) have an increased risk of developing NIDDM.

Preferably such a DNA sequence is one containing a mutation of G to A in the third position of codon 326 of the regulatory p85 α subunit of PI3K gene such that methionine³²⁶
30 is substituted by isoleucine. This mutation is at a domain boundary, directly in front of an SH2 domain of p85 α . The region is located downstream of a hydrophobic region the function of which is not known but which might be membrane-associated, as the subunit

SUBSTITUTE SHEET (RULE 26)

apparently acts as an adaptor allowing the kinase to associate with the membrane. Figure 1 is a diagram of the PI3K p85 α that shows the position of the codon326 mutation relative to an SH2 domain.

5 Apart from the missense mutations leading to substitution of one or more amino acids in the regulatory p85 α subunit of PI3K protein, it is expected that silent mutations in the form of silent single nucleotide substitutions may have an influence on splicing (if it is located within a few bases of the exon-intron boundary), mRNA stability (if it changes the folding or the accessibility of the mRNA to RNAses), or protein stability (since codons may be
10 changed into rare codons which may be translated more slowly and thereby influence the folding/stability of the protein (cf. Ikemura, T. and Ozeki, H.: Codon usage and transfer RNA contents: organism-specific codon-choice patterns in preference to the isoacceptor contents. *Cold Spring Harbor Symp. Quant. Biol.* 47:1087-1096, 1982; Grosjean, F. and Fiers, W.: Preferential codon usage in prokaryotic genes: the optimal codon-anticodon
15 interaction energy and the selective codon usage in efficiently expressed genes. *Gene* 18:199-209, 1982). Such silent mutations may also be correlated with other important genetic changes and may therefore be useful as markers for detection of important genetic variants.

20 The length of the nucleic acid sequence of the invention, such as a cDNA isolate, may vary widely depending on the intended use. For use as an oligonucleotide probe for hybridisation purposes, the cDNA segment may be as short as 17 nucleotides. For expression in a living system as defined above, the cDNA isolate will typically comprise the full-length cDNA sequence encoding the regulatory p85 α subunit of PI3K.

25

The nucleic acid sequence of the invention comprising the mutation in the cDNA sequence encoding the regulatory p85 α subunit of PI3K may suitably be of genomic or cDNA origin, for instance obtained by preparing a genomic or cDNA library and screening for DNA sequences coding for all or part of the regulatory p85 α subunit of PI3K by hybridization
30 using synthetic oligonucleotide probes in accordance with standard techniques (cf. Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989). The probes used should be specific for the mutation. Alternatively, the DNA

sequence encoding wild-type of the regulatory p85 α subunit of PI3K may be modified by site-directed mutagenesis using synthetic oligonucleotides containing the mutation for homologous recombination in accordance with well-known procedures.

5 The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202, Saiki et al., Science 239, 1988, pp. 487-491, or PCR Protocols, 1990, Academic Press, San Diego, California, USA.

The DNA isolate of the invention may also be prepared synthetically by established
10 standard methods, e.g. the phosphoamidite method described by Beaucage and Caruthers, Tetrahedron Letters 22 (1981), 1859 - 1869, or the method described by Matthes et al., EMBO Journal 3 (1984), 801 - 805. According to the phosphoamidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed and ligated. This procedure may preferably be used to prepare shorter segments of the
15 regulatory p85 α subunit of PI3K encoding DNA sequence.

The recombinant vector into which the DNA isolate is inserted may be any vector which may conveniently be subjected to recombinant DNA procedures. The choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an
20 autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated (e.g. a viral vector).

25

The recombinant vector is preferably an expression vector in which the DNA sequence encoding the regulatory p85 α subunit of PI3K mutant is operably connected to additional segments required for transcription of the DNA. In general, the vector is derived from plasmid or viral DNA or may contain elements of both. The term "operably linked"
30 indicates that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in a promoter and proceeds through the DNA sequence coding for the regulatory p85 α subunit of PI3K.

The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the mutant DNA encoding the regulatory p85 α subunit of PI3K in mammalian cells are the SV40 promoter (Subramani et al., Mol. Cell Biol. 1 (1981), 854 -864), the MT-1 (metallothionein gene) promoter (Palmiter et al., Science 222 (1983), 809 - 814) or the adenovirus 2 major late promoter.

- 10 The mutant DNA sequence encoding the regulatory p85 α subunit of PI3K may also be operably connected to a suitable terminator, such as the human growth hormone terminator (Palmiter et al., op. cit.). The vector may further comprise elements such as polyadenylation signals (e.g. from SV40 or the adenovirus 5 Elb region), transcriptional enhancer sequences (e.g. the SV40 enhancer) and translational enhancer sequences (e.g. the
15 ones encoding adenovirus VA RNAs).

The recombinant vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. An example of such a sequence is the SV40 origin of replication. The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the gene coding for dihydrofolate reductase (DHFR) or one which confers resistance to a drug, e.g. neomycin, hygromycin or methotrexate.

The procedures used to ligate the DNA sequences coding for the regulatory p85 α subunit of PI3K, the promoter and the terminator, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al., op.cit.).

In a further aspect, the present invention relates to a variant of the regulatory p85 α subunit of PI3K containing at least one amino acid substitution, in particular a variant containing at least one amino acid substitution at a site where the substitution interferes with the binding of the regulatory p85 α subunit of PI3K to insulin receptor kinase or insulin signalling

proteins, e.g. IRS-1 and/or IRS-2, or a fragment thereof including said substitution. Examples of such variants are variants in which methionine³²⁶ is substituted by isoleucine.

The living system into which the DNA isolate of the invention is introduced may be a cell 5 which is capable of producing the regulatory p85 α subunit of PI3K and which has the appropriate signal transduction pathways. The cell is preferably a eukaryotic cell, such as a vertebrate cell, e.g. a *Xenopus laevis* oocyte or mammalian cell, in particular a mammalian cell. Examples of suitable mammalian cell lines are the COS (ATCC CRL 1650), BHK (ATCC CRL 1632, ATCC CCL 10), CHL (ATCC CCL39) or CHO (ATCC CCL 61) cell 10 lines. Methods of transfecting mammalian cells and expressing DNA sequences introduced in the cells are described in e.g. Kaufman and Sharp, J. Mol. Biol. 159 (1982), 601 - 621; Southern and Berg, J. Mol. Appl. Genet. 1 (1982), 327 - 341; Loyter et al., Proc. Natl. Acad. Sci. USA 79 (1982), 422 - 426; Wigler et al., Cell 14 (1978), 725; Corsaro and Pearson, Somatic Cell Genetics 7 (1981), 603, Graham and van der Eb, Virology 52 15 (1973), 456; and Neumann et al., EMBO J. 1 (1982), 841 - 845.

The mutant DNA sequence encoding the regulatory p85 α subunit of PI3K may then be expressed by culturing a cell as described above in a suitable nutrient medium under conditions which are conducive to the expression of the regulatory p85 α subunit of PI3K- 20 coding DNA sequence. The medium used to culture the cells may be any conventional medium suitable for growing mammalian cells, such as a serum-containing or serum-free medium containing appropriate supplements. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. in catalogues of the American Type Culture Collection).

25

The living system according to the invention may also comprise a transgenic animal. A transgenic animal is one in whose genome a heterologous DNA sequence has been introduced. In particular, the transgenic animal is a transgenic non-human mammal, mammals being generally provided with appropriate signal transduction pathways. For the 30 present purpose, it is generally preferred to employ smaller mammals, e.g. rabbits or rodents such as mice or rats.

For expression of the mutant regulatory p85 α subunit of PI3K gene of the invention in transgenic animals, a DNA sequence encoding the mutant regulatory p85 α subunit of PI3K is operably linked to additional DNA sequences required for its expression to produce expression units. Such additional sequences include a promoter as indicated above, as well as sequences providing for termination of transcription and polyadenylation of mRNA. Construction of the expression unit for use in transgenic animals may conveniently be done by inserting a mutant DNA sequence encoding the regulatory p85 α subunit of PI3K into a vector containing the additional DNA sequences, although the expression unit may be constructed by essentially any sequence of ligations.

10

The expression unit is then introduced into fertilized ova or early-stage embryos of the selected host species. Introduction of heterologous DNA may be carried out in a number of ways, including microinjection (cf. US 4,873,191), retroviral infection (cf. Jaenisch, Science 240, 1988, pp. 1468-1474) or site-directed integration using embryonic stem cells (reviewed by Bradley et al., Bio/Technology 10, 1992, pp. 534-539). The ova are then implanted into the oviducts or uteri of pseudopregnant females and allowed to develop to term. Offspring carrying the introduced DNA in their germ line can pass the DNA on to their progeny, allowing the development of transgenic populations.

20 General procedures for producing transgenic animals are known in the art, cf. for instance, Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory, 1986; Simons et al., Bio/Technology 6, 1988, pp. 179-183; Wall et al., Biol. Reprod. 32, 1985, pp. 645-651; Buhler et al., Bio/Technology 8, 1990, pp. 140-143; Ebert et al., Bio/Technology 6: 179-183, 1988; Krimpenfort et al., Bio/Technology 9: 844-25 847, 1991, Wall et al., J. Cell. Biochem. 49: 113-120, 1992; US 4,873,191, US 4,873,316; WO 88/00239, WO 90/05188; WO 92/11757 and GB 87/00458. Techniques for introducing heterologous DNA sequences into mammals and their germ cells were originally developed in the mouse. See, e.g. Gordon et al., Proc. Natl. Acad. Sci. USA 77: 7380-7384, 1980, Gordon and Ruddle, Science 214: 1244-1246, 1981; Palmiter and 30 Brinster, Cell 41: 343-345, 1985; Brinster et al., Proc. Natl. Acad. Sci. USA 82: 4438-4442, 1985; and Hogan et al. (ibid.). In brief, in the most efficient route used to date in the generation of transgenic mice, several hundred linear molecules of the DNA of interest are

injected into one of the pro-nuclei of a fertilized egg according to techniques which have become standard in the art. Injection of DNA into the cytoplasm of a zygote can also be employed. Similar procedures may be employed to produce transgenic individuals of other species.

5

In one embodiment of the present method of detecting the presence of a mutation in the regulatory p85 α subunit of PI3K gene, a biological sample is obtained from a subject, DNA (in particular genomic DNA) is isolated from the sample and digested with a restriction endonuclease which cleaves DNA at the site of the mutation, and cleavage of the DNA within the gene encoding the regulatory p85 α subunit of PI3K at this site is determined. After digestion, the resulting DNA fragments may be subjected to electrophoresis on an agarose gel. DNA from the gel may then be blotted onto a nitrocellulose filter and hybridised with a radiolabelled probe. The probe may conveniently contain a DNA fragment of the regulatory p85 α subunit of PI3K gene spanning the mutation (substantially according to the method of E.M. Southern, J. Mol. Biol. **98**, 1975, pp. 503, e.g. as described by B.J. Conner et al., Proc. Natl. Acad. Sci. USA **80**, 1983, pp. 278-282).

In a variant of this embodiment, the DNA isolated from the sample may be amplified prior to digestion with the restriction endonuclease. Amplification may suitably be performed by polymerase chain reaction (PCR) using oligonucleotide primers based on the appropriate sequence of the regulatory p85 α subunit of PI3K spanning the site(s) of mutation, essentially as described by Saiki et al., Science **230**, 1985, pp. 1350-1354. After amplification, the amplified DNA may be digested with the appropriate restriction endonuclease and subjected to agarose gel electrophoresis. The restriction pattern obtained may be analysed, e.g. by staining with ethidium bromide and visualising bands in the gel by means of UV light. As a control, wild-type DNA encoding the regulatory p85 α subunit of PI3K (i.e. not containing the mutation) may be subjected to the same procedure, and the restriction patterns may be compared.

In the method of the invention, the sample is preferably analysed for a mutation located at a site where amino acid substitution interferes with the binding of the regulatory p85 α subunit of PI3K to insulin receptor kinase or insulin signalling proteins, e.g. IRS-1 and /or

SUBSTITUTE SHEET (RULE 26)

IRS-2. We have found that such a mutation is the mutation of G to A in the third position of codon 326 of the gene encoding the regulatory p85 α subunit of PI3K.

A further embodiment of the method of the invention is an adaptation of the method 5 described by U. Landegren et al., Science 241, 1988, pp. 1077-1080, which involves the ligation of adjacent oligonucleotides on a complementary target DNA molecule. Ligation will occur at the junction of the two oligonucleotides if the nucleotides are correctly base paired.

10 In a still further embodiment of the present method, the DNA isolated from the sample may be amplified using oligonucleotide primers corresponding to segments of the gene coding for the regulatory p85 α subunit of PI3K. The amplified DNA may then be analysed by hybridisation with a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least part of the gene encoding the regulatory p85 α subunit of PI3K and
15 containing a mutation of at least one nucleotide, which mutation corresponds to the mutation the presence of which in the gene encoding the regulatory p85 α subunit of PI3K is to be detected. As a control, the amplified DNA may furthermore be hybridised with a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least part of the wild-type gene encoding the regulatory p85 α subunit of PI3K. This procedure is, for
20 instance, described by DiLella et al., Lancet 1, 1988, pp. 497-499. Other PCR-based methods which may be used in the present invention are the allele-specific PCR method described by R. Saiki et al., Nature 324, 1986, pp. 163-166, or D.Y. Wu et al., Proc. Natl. Acad. Sci. USA 86, 1989, pp. 2757-2760, which uses primers specific for the mutation in the regulatory p85 α subunit of PI3K gene; the ligase chain reaction; restriction
25 generating PCR; oligo nucleotide ligation assay; mini-sequencing; primer extension; and the heteroduplex method, cf. Landegren, U. (ed.) Laboratory Protocols for Mutation Detection. Oxford University Press, 1996.

Other methods of detecting mutations in DNA are reviewed in U. Landegren, GATA 2,
30 1992, pp. 3-8. A currently preferred method of detecting mutations is by single stranded conformation polymorphism (SSCP) analysis substantially as described by Orita M, Iwahana H, Kanazawa H, Hayashi K, Sekiya T: Detection of polymorphisms of human DNA by gel

electrophoresis as single stranded conformational polymorphisms. *Proc Natl Acad Sci* 86:2766-70, 1989, or Orita et al., *Genomics* 5, 1989, pp. 874-879.

The label substance with which the probe is labelled is preferably selected from the group 5 consisting of enzymes, coloured or fluorescent substances, or radioactive isotopes.

Examples of enzymes useful as label substances are peroxidases (such as horseradish peroxidase), phosphatases (such as acid or alkaline phosphatase), β -galactosidase, urease, glucose oxidase, carbonic anhydrase, acetylcholinesterase, glucoamylase, lysozyme, malate 10 dehydrogenase, glucose-6-phosphate dehydrogenase, β -glucosidase, proteases, pyruvate decarboxylase, esterases, luciferase, etc.

Enzymes are not in themselves detectable but must be combined with a substrate to catalyse a reaction the end product of which is detectable. Examples of substrates which may be em- 15 ployed in the method according to the invention include hydrogen peroxide/tetramethylbenzidine or chloronaphthole or o-phenylenediamine or 3-(p-hydroxyphenyl) propionic acid or luminol, indoxyl phosphate, p-nitrophenylphosphate, nitrophenyl galactose, 4-methyl umbelliferyl-D-galactopyranoside, or luciferin.

20 Alternatively, the label substance may comprise coloured or fluorescent substances, including gold particles, coloured or fluorescent latex particles, dye particles, fluorescein, phycoerythrin or phycocyanin.

In a particularly favoured embodiment, the probe is labelled with a radioactive isotope. 25 Radioactive isotopes which may be used for the present purpose may be selected from I-125, I-131, In-111, H-3, P-32, C-14 or S-35. The radioactivity emitted by these isotopes may be measured in a beta- or gamma-counter or by a scintillation camera in a manner known *per se*.

30 For use in the present method, the invention further relates to a test kit for detecting the presence of a mutation in the gene encoding the regulatory p85 α subunit of PI3K, the kit comprising

- (a) a restriction endonuclease which cleaves DNA at the site of the mutation,
- (b) a first DNA sequence corresponding to at least part of the wild-type gene encoding the 5 regulatory p85 α subunit of PI3K and/or
- (c) a second DNA sequence corresponding to at least part of the gene encoding the regulatory p85 α subunit of PI3K and containing a mutation of at least one nucleotide, which mutation corresponds to the mutation the presence of which in the gene encoding the 10 regulatory p85 α subunit of PI3K is to be detected. Said mutation is preferably the codon326 mutation resulting in a methionin to isoleucin substitution.

The first DNA sequence may, for instance, be obtained from genomic DNA or cDNA encoding the regulatory p85 α subunit of PI3K obtained from a healthy subject (a non-15 mutation carrier). The second DNA sequence may conveniently be a DNA construct according to the invention.

For use in the present method, the invention further relates to a test kit for detecting the presence of a mutation in the gene encoding the regulatory p85 α subunit of PI3K, the kit 20 comprising

- (a) means for amplifying DNA, and
- (b) a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least 25 part of the gene encoding the regulatory p85 α subunit of PI3K and containing a mutation of at least one nucleotide, which mutation corresponds to the mutation the presence of which in the gene encoding the regulatory p85 α subunit of PI3K is to be detected. Preferably, the mutation the presence of which is to be detected is the codon 326 methionin to isoleucin mutation of p85 α .

30

Appropriate means for amplifying DNA (typically genomic DNA isolated from the biological sample) include, for instance, oligonucleotide primers, appropriate buffers and a

thermostable DNA polymerase.

The kit may further comprise a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least part of the wild-type gene encoding the regulatory p85 α subunit of 5 PI3K.

The invention also relates to the use of a nucleic acid sequence of the invention or a segment thereof as a probe, preferably an oligonucleotide probe, for detection of the presence of a mutation in a regulatory subunit, preferably the p85 α subunit or any isoform 10 or homologous polypeptide thereof, of PI3K, the regulatory subunit being preferably isolated from a biological sample obtained from a subject.

The invention is further illustrated in the following example which is not intended in any way to limit the scope of the invention as claimed.

15

EXAMPLE

Subjects. For studies of intravenous glucose and tolbutamide stimulated serum insulin and C-peptide release 380 study participants were randomly recruited from a population of 20 young individuals aged 18-32 years, who in 1979/80 and again in 1984/85 as children had participated in blood pressure surveys in a representative and specified part of Copenhagen City. All were Danish Caucasians by self-identification. Physiological characteristics and other genetic analyses obtained in the present population sample have been presented previously, cf. Clausen et al. The Lancet 346:397-402, 1995. Prior to the participation in 25 the study informed consent was obtained for all subjects. The study was approved by the Ethical Committee of Copenhagen and was in accordance with the principles of the Declaration of Helsinki.

Measurements of glucose disappearance constant (Kg), insulin sensitivity (Si) and 30 glucose effectiveness (Sg). Each subject underwent an intravenous glucose tolerance test (IVGTT) after a 12 h overnight fasting period. Baseline values of serum insulin, serum C-peptide and plasma glucose were taken in duplicate with 5 min intervals. Glucose was

injected i.v. in the contralateral antecubital vein over a period of 1 min (0.3 g/kg body weight of 50% glucose). At 20 min following the end of the glucose injection, a bolus of 3 mg tolbutamide/kg body weight (Rastinon, Hoechst, Germany) was injected during 5 sec to elicit a secondary pancreatic beta cell response. Venous blood was sampled at 2, 4, 8, 19, 22, 30, 40, 50, 70, 90, 180 min, timed from the end of the glucose injection for measurements of plasma glucose, serum insulin and serum C-peptide. All the IVGTT's were done by the same investigator. Glucose induced acute serum insulin and C-peptide responses (0-8 min) were calculated by means of the trapezoidal rule as the incremental values (area under the curve when expressed above basal values). Insulin sensitivity index (Si) and glucose effectiveness (Sg) were calculated using the Bergman MINMOD computer program developed specifically for the combined intravenous glucose and tolbutamide tolerance test, cf. Clausen et al. op cit. Intravenous glucose disappearance constant (Kg) was calculated as the slope of the line relating the natural logarithm of the glucose concentration to the time between 8 and 19 minutes after the glucose bolus. Plasma concentrations of glucose, serum specific insulin and serum C-peptide were measured as described.

Chemical analysis. Chemical analyses were performed as described previously (cf. Almind K, Bjørbæk C, Vestergaard H, Hansen T, Echwald S, Pedersen O: Amino acid polymorphisms of insulin receptor substrate-1 in non-insulin-dependent diabetes mellitus. *The Lancet* 342:828-32, 1993). Briefly, measurements of glucose in plasma was done by a hexokinase method and serum C-peptide concentrations were analyzed by RIA.

Isolation of genomic DNA from whole blood. Genomic DNA was isolated from human leucocyte nuclei isolated from whole blood by digestion with proteinase K followed by phenol extraction on an Applied Biosystems 341 Nucleic Acid Purification System (Applied Biosystems Inc., Foster City, California, USA)

Muscle biopsies. Needle biopsies of the vastus lateralis muscle (500 mg) from subjects were obtained during local anesthesia (1 % Lidocaine, without epinephrine). A modified Bergström needle was used (Stille-Werner, Copenhagen Denmark), and samples were taken from about 20 cm above the knee. Biopsies were blotted to remove blood and connective

tissue and frozen in liquid nitrogen within 30 seconds.

Preparation of total RNA from skeletal muscle. Muscle biopsies were thawed and homogenized in a 4 M guanidinium thiocyanate solution. Total RNA was isolated using a 5 394 RNA/DNA extractor (Applied Biosystems Inc., Foster City, California, USA). Purity and concentration of the samples were determined by measuring the 260/280 nm absorption ratio.

cDNA synthesis. cDNA was synthesized in volumes of 25 μ l containing (in final 10 concentrations) 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3mM $MgCl_2$, 10 mM DDT, 0.2 mM dNTP's, 40 U Rnasin (Promega, Madison, WI), 0.625 μ g Oligo (dT)₁₈, 400 U Molony murine leukemia virus reverse transcriptase (M-MLV RT) (Life Technologies, Grand Island, NY), and 1.0 μ g of total muscle RNA. The reactions were performed at 37°C for 1 h, followed by enzyme inactivation for 10 min at 95°C. Samples were diluted to 15 a final volume of 200 μ l and stored at -20°C.

Oligonucleotides used for amplification. Oligonucleotides, 18-21 mers in length and with one or two G's and/or C's at the 3' end, were synthesized on an Applied Biosystems 394 DNA/RNA Synthesizer and applied to a NABTM -10 column (Pharmacia P-L Biochemicals 20 Inc., Milwaukee, WI) and used without further purification. Biotin labelling of primers was done during synthesis using DMT-Biotin-C6-PA (Genosys Biotechnologies Inc., Great Britain). The human p85 α has been published by Skolnik (Skolnik et al., 1991) (GenBank, Genetics Computer Group (Madison, WI), accession number M61906). The published amino acid sequence has 97% homology to bovine p85 α cDNA (M61745), 95% to mouse 25 p85 α cDNA (M60651) and only 59% to bovine p85 β cDNA (M61746). Primers for polymerase chain reaction (PCR) amplification of human p85 α cDNA were designed from areas of the cDNA sequence, with low homology to bovine p85 β (Fig. 2).

PCR amplification of cDNA. A primary PCR amplification was carried out with 5 μ l 30 cDNA as template. The assay conditions were: 10 mM Tris-HCl (pH 9.0), 50 mM KCl, 1.5 mM $MgCl_2$, 0.1% Triton X-100, 0.2 mM dNTP's, 0.2 μ M of each oligonucleotide primer, and 0.6 U Taq DNA Polymerase (Promega, Madison, WI) (all in final

concentrations and total volume of 25 μ l). The mixture was overlaid with mineral oil and after initial denaturation at 95°C for 4 min, the samples were subjected to 35 cycles of amplification: denaturation at 95°C for 1 min, annealing at 55°C for 1 min and extension at 72°C for 2 min (Omnigene Thermocycler, Hybaid Limited, Middlesex, U.K.). A final step 5 with 5 min extension at 72°C was performed to ensure full-length segments.

Primary SSCP screening for variants in the regulatory p85 α subunit of PI3K gene. The coding region of PI3K p85 α subunit was scanned by PCR-SSCP at the nucleotide level. 10 cDNA from 70 NIDDM patients and 12 healthy control subjects without any known family history of NIDDM was included in the primary SSCP screening. SSCP-analysis (according to the method of Orita et al., Genomics 5, 1989, pp. 874-879) was performed on DNA-segments of 250-300 bp with an overlap of 60 base pairs between segments. DNA-segments for SSCP analysis were PCR-amplified in a standard PCR-reaction, using specific primers 15 as described above. PCR-reactions were performed as described above and 3 μ l of the PCR-reaction was mixed with 12 μ l of SSCP loading buffer (95 % deionised formamide, 5 % NaOH, xylene blue, bromo-phenol blue), and 2 μ l were analyzed on a 38 x 31 x 0.04 cm, non-denaturing 5 % acrylamide (49:1 acrylamide:bisacrylamide) gel in adjacent lanes. For heteroduplex formation samples were loaded after heating at 95°C for 5 min. and 20 subsequently incubating 15 min. on ice. Each DNA-segment was analyzed at two different gel conditions. A "cold gel", containing 1 % glycerol, 1 X TBE-buffer (90 mM Tris-Borate, 2.5 mM EDTA), and 5 % acrylamide (49:1, acrylamide:bisacrylamide) was run at 8 °C in a cold-room for 3 to 5 h with an applied power of 35 watt. A "warm" gel containing 5 % glycerol, 1 X TBE-buffer (90 mM Tris-Borate, 2.5 mM EDTA), and 5 % 25 acrylamide (49:1, acrylamide:bisacrylamide), was run at 24°C for 3 to 4 h, applying 65 watt. The electrophoresis buffer was 1/2 x TBE (45 mM Tris Base, 1.25 mM EDTA). The power supplies used were equipped with a temperature controlling device, to assure constant running temperature (Temperature Controller, Stratagene, La Jolla, CA, USA). After electrophoresis, the gels were transferred to Whatman 3 MM filter paper, covered in 30 plastic wrap, and autoradiografed at -80°C, for 3 h to 3 days using intensifying screens. Autoradiografies were analyzed visually to reveal variations in migration. SSCP screening and subsequent direct sequencing revealed 4 nucleotide variants, shown in Table 1 below.

SSCP SEGMENT	SET OF PRIMERS	NUCLEOTIDE CHANGE POSITION/BASE	AMINO ACID CHANGE CODON/A.A.	GENOTYPE FREQUENCY NIDDM		
				Control		
2.	3-4	261/C → T	73/Tyr → Tyr	ho	9	0
				he	29	4
				wt	32	8
4.	7-8	663/T → G	207/Ile → Ile	ho	0	0
				he	3	1
				wt	67	11
4.	7-8	810/A → G	256/Lys → Lys	ho	0	0
				he	2	0
				wt	68	12
5.	9-10	1020/G → A	326/Met→ Met	ho	1	0
				he	5	1
				wt	64	11

Table 1. Localisation, character and genotype frequency of nucleotide variants in PI3K 5 p85 α subunit. The base number is calculated from first sequenced base (accession number M69106), the codon number is calculated from first coding methionin (code 1, base number 43-45, sequence accession number M69106).

Primers used for PCR amplification of the codon 326 polymorphism for SSCP are listed in 10 table 2. Figure 2 shows the distribution of primers, primary and secondary segments in SSCP scanning of p85 α . The numbers refer to the primer numbers in Table 2, e.g. (1-8) refers to the primers PI 1 through PI8 of Table 2.

NAME	SPEC	GENE	SEQUENCE	BASE	CONC.
	(BIO)		5' - 3'	NUMBER	PMOL/ul
PI 1		PI3K, P85	caa cca ggc tca act gtt gc	1 to 22	397
PI 2		PI3K, P85	tac gtt ccc gga aag tcc cc	240 to 60	402
PI 3		PI3K, P85	gaa att ggc tgg tta aat ggc	195 to 216	494
PI 4		PI3K, P85	gag ctt gat aag aag agg cg	437 to 447	359
PI 5		PI3K, P85	gat ctt gca gag cag ttg gc	390 to 410	496
PI 6		PI3K, P85	acg gct gct gga atg aca gg	628 to 647	407
PI 7		PI3K, P85	tgg ctg acg ctt tca aac gc	586 to 606	528
PI 8		PI3K, P85	gaa tct gaa aag cat agg gc	848 to 867	425
PI 9		PI3K, P85	ag ctc tct caa acc tcc agc	787 to 807	695
PI 10		PI3K, P85	cga gat atc tcc cca gta cc	1139 to 1157	757
PI 11		PI3K, P85	ac tac tgt agc caa caa cgg	984 to 1104	638
PI 12		PI3K, P85	gtc aga gaa gcc ata ttg	1203 to 1224	402
PI 13		PI3K, P85	t tac act aag gaa agg ggg	1143 to 1169	363
PI 14		PI3K, P85	ga ttg ctt ggg atg tgc ggg	1454 to 1473	337
PI 15		PI3K, P85	cac tca gtt tca aga aaa aag ta	1401 to 1424	629
PI 16		PI3K, P85	gc ttc ttc aag tct tct tcc	1677 to 1696	394
PI 17		PI3K, P85	tga taa gtt gaa gtc tgc	1627 to 1645	366
PI 18		PI3K, P85	a tct tca tca tct tcc acc	1870 to 1879	230
PI 19		PI3K, P85	ag ttg aac gag tgg ttg ggc	1817 to 1836	418
PI 20		PI3K, P85	g aaa gcc ata gcc agt tgc	2068 to 2086	291
PI 21		PI3K, P85	gg tgg acg gcg aag taa agc	2029 to 2048	377
PI 22		PI3K, P85	agc cct ttg ctt tcc aga gg	2264 to 2283	335

Table 2: Nucleotide sequences of DNA primers used for PCR amplification to detect the regulatory p85 α subunit of PI3K amino acid substitution at codon 326 by SSCP gel 20 analysis.

Secondary screening for additional codon 326 mutation carriers. Additionally, in a random sample of unrelated, healthy young volunteers in whom an intravenous glucose tolerance test in combination with tolbutamide injection was performed, 263 wild-type 25 (Wt), 109 heterozygous (He) and 8 homozygous (Ho) individuals were identified. Glucose disappearance constant (Kg) and glucose effectiveness (Sg) were significantly different among the groups (Kg: Wt $2.3 \pm 1.1 \times 10^{-2} \text{ min}^{-1}$, He 2.3 ± 1.0 , Ho 1.4 ± 0.6 (mean \pm SD)($p=0.02$); Sg: Wt $2.1 \pm 0.7 \times 10^{-2} \text{ min}^{-1}$, He 2.2 ± 0.6 , Ho 1.7 ± 0.6 ($p=0.03$)). No significant difference was observed in insulin sensitivity (Wt $15.3 \pm 9.5 \times 10^{-3} \text{ L min}^{-1}$ 30 pmol^{-1} , He 15.1 ± 8.6 , Ho 10.3 ± 8.3 ($p=0.20$)). Both Kg and Sg were significantly reduced among Ho subjects compared to the combined values for Wt and He carriers ($p=0.004$ and $p=0.03$, respectively). Moreover, in Ho carriers a 30% reduction in insulin sensitivity

($p=0.08$) was found. All homozygous subjects were validated using direct sequencing. It was concluded that in homozygous carriers the common codon 326^{Met→Ile} variant of the p85 α subunit of PI3K is associated with a decreased glucose disappearance constant possibly resulting in intravenous glucose tolerance and a decreased glucose effectiveness in 5 young, healthy individuals.

Allele specific oligonucleotide hybridization: Genomic DNA (100 ng) was PCR amplified with primers 10 and 11 (Fig. 2) as described above except for addition of 2.0 mM MgCl₂ and 10% DMSO. The samples were subjected to 40 cycles of amplification: denaturation at 10 95 °C for 30 sec, annealing at 54 °C for 30 sec, extension at 72 °C for 30 sec and final extension for 5 min at 72 °C using GeneAmp PCR system 9600 (Perkin Elmer, Norwalk, CT). PCR samples (15 μ l) were denatured in 85 μ l 0.2 M NaOH, 1 M NaCl and 2 x 45 μ l were transferred to Hybond-N⁺ filters (Amersham, Buckinghamshire, Great Britain).

15 The membranes were crosslinked twice in a UV Stratalink (Stratagene, La Jolla, CA) at 1200 x 100 μ Joule. Membranes were prehybridized for 1 hour in 6xSSPE, 5xDenhardt, 0.1% SDS and 100 μ g/ml denatured salmon sperm. Hybridization was performed with allele specific probes (wild type: 5' AAC AAT ATG TCC TTA CA; variant 5' AAC AAT ATA TCC TTA CA) for 3 hours in 6xSSPE, 5xDenhardt, 0.1% SDS. Probes were 5'-end-
20 labeled by T4-kinase (Promega, Madison, WI) as described (ref) using (Y-³²P) ATP (Amersham, Buckinghamshire, Great Britain). After hybridization membranes were washed for 2x0.5 h in 6xSSPE and 0.1% SDS at 45 °C for wild type probe and 41 °C for variant probe. Filters were applied to PhosphorImager Screens for 2-10 h and visualized on an ImageQuant, PhosphorImager available from Molecular Dynamics.

25

Direct sequencing. SSCP-variants were sequenced using either an Automated Laser Fluorescence Sequencer (LKB Pharmacia Uppsala, Sweden) and using Fluore-dATP (Pharmacia/Boeringer Mannheim) as label in the AutoRead Sequencing Kit (Pharmacia, Uppsala, Sweden), or by standard dideoxynucleotide sequencing using the Sequenase 2.0
30 (United States Biochemical Corporation, Cleveland, OH) and (α -S³⁵) dATP (Amersham International plc, Buckinghamshire, Great Britain) according to protocols supplied by the producer.

DNA for sequencing was PCR-amplified using 5 pmol/100 μ l reaction of one biotinylated primer, and single-stranded DNA was isolated using streptavidin-coated magnetic beads (Dynabeads, Oslo, Norway). DNA was sequenced as single stranded DNA according to 5 protocols, using internal sequencing primers.

Statistics. Chi-square analysis was applied to test for significance of differences in allele and genotype frequencies. A non-parametric Kruskal-Wallis test or a Mann-Whitney test was used for comparison between groups. Data are means \pm SEM. A p-value < 0.05 (two-tailed) was considered significant. Statistical Package of Social Science (SPSS) for Windows, version 6.01, was used for statistical analysis.

CLAIMS

1. A nucleic acid sequence comprising a DNA sequence encoding a regulatory subunit of phosphatidylinositol 3-kinase (PI3K), the DNA sequence containing a mutation of at least
5 one nucleotide, or comprising a segment of the DNA sequence including said mutation for use as a diagnostic tool, a marker or a probe.
2. A nucleic acid sequence according to claim 1, wherein the regulatory subunit is the p85 α subunit of human PI3K.
- 10 3. A nucleic acid sequence according to claim 1 or 2, wherein the mutation is a mutation of
C to T in nucleotide 261;
T to G in nucleotide 663;
A to G in nucleotide 810;
15 G to A in nucleotide 1020.
4. A nucleic acid sequence according to any one of claims 1, 2 or 3, wherein the mutation gives rise to at least one amino acid mutation.
- 20 5. A nucleic acid sequence according to claim 4, wherein the mutation is G to A in the third position of codon 326 such that methionine³²⁶ is substituted by isoleucine.
6. A recombinant vector comprising a nucleic acid sequence according to any one of claims 1-5.
- 25 7. A living system containing a nucleic acid sequence according to any one of claims 1-5 or a recombinant vector according to claim 6 and capable of expressing a regulatory subunit of PI3K wherein at least one amino acid is substituted.
- 30 8. A living system according to claim 7 which comprises a mammalian cell line or a transgenic animal, in particular a transgenic non-human mammal.

9. A method of detecting the presence of a mutation in a gene encoding a regulatory subunit of PI3K, the method comprising obtaining a biological sample from a subject and analysing the sample for the presence of a nucleic acid sequence according to any one of claims 1-5.
- 5 10. A method according to claim 9, wherein the regulatory subunit is p85 α of human PI3K.
11. A method according to any one of claims 9 or 10, wherein the sample is analysed for a mutation of
- C to T in nucleotide 261,
- 10 T to G in nucleotide 663,
- A to G in nucleotide 810, or
- G to A in nucleotide 1020
- of the p85 α regulatory subunit of PI3K.
- 15 12. A method according to claim 11, wherein the sample is analysed for a mutation of G to A in the third position of codon 326 giving rise to substitution of methionine³²⁶ by isoleucine.
13. A method according to any one of claims 9 to 12, wherein a biological sample is
- 20 obtained from a subject, DNA is isolated from the sample and digested with a restriction endonuclease which cleaves DNA at the site of the mutation, and cleavage of the DNA within the gene encoding the p85 α regulatory subunit of PI3K at this site is determined.
14. A method according to claim 13, wherein the restriction pattern of the DNA after
- 25 digestion with the restriction endonuclease is compared to the restriction pattern obtained with a negative control comprising at least a portion of wild-type DNA encoding the p85 α regulatory subunit of PI3K.
15. A method according to claim 13, wherein the restriction pattern of the DNA after
- 30 digestion with the restriction endonuclease is compared to the restriction pattern obtained with a positive control comprising at least a portion of DNA encoding the p85 α regulatory subunit of PI3K and containing the mutation.

16. A method according to claim 15, wherein the positive control comprises a nucleic acid sequence according to any one of claims 1-5.

5 17. A method according to claim 13, which further comprises amplifying the DNA isolated from the sample prior to digestion with the restriction endonuclease.

18. A method according to any one of claims 9-17 for determining predisposition to impaired glucose tolerance in a subject.

10

19. A method according to the previous claim, wherein the impaired glucose tolerance is related to an impaired glucose disappearance constant in a subject.

20. A method according to any of claims 9-17 for determining predisposition to decreased
15 glucose effectiveness.

21. A method according to any one of claims 9-17 for determining predisposition to decreased insulin sensitivity in a subject.

20 22. A method according to any one of claims 9 or 10, wherein a biological sample is obtained from a subject, DNA is isolated from the sample, the DNA is amplified and hybridised to a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least part of the gene encoding a regulatory subunit of PI3K and containing a mutation of at least one nucleotide, which mutation corresponds to the mutation the presence of
25 which in said gene is to be detected, and hybridisation of the probe to the DNA is determined.

23. A method according to claim 22, wherein the amplified DNA is further hybridised to a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least part of
30 the wild-type gene encoding a regulatory subunit of PI3K, and hybridisation of the probe to the DNA is determined.

25

24. A method according to claim 22 or 23, wherein the label substance with which the oligonucleotide probe containing the mutation is labelled is different from the label substance with which the oligonucleotide probe corresponding to the wild-type DNA is labelled.

5

25. A method according to any one of claims 22 or 23, wherein the label is selected from the group consisting of enzymes, coloured or fluorescent substances, and radioactive isotopes.

10 26. A method according to any of claims 22-25 for determining predisposition to impaired glucose tolerance in a subject.

27. A method according to any one of claims 22-25 wherein the impaired glucose tolerance is related to an impaired glucose disappearance constant in a subject.

15

28. A method according to any of claims 22-25 for determining predisposition to decreased glucose effectiveness.

29. A method according to any one of claims 22-25 for determining predisposition to
20 decreased insulin sensitivity in a subject.

30. A diagnostic composition for detecting the presence of a mutation in the gene encoding a regulatory subunit of PI3K, the composition comprising a nucleic acid sequence according
25 to any one of claims 1-5.

31. A test kit for detecting the presence of a mutation in the gene encoding a regulatory subunit of PI3K, the kit comprising

30 (a) a restriction endonuclease which cleaves DNA at the site of the mutation,

(b) a first DNA sequence corresponding to at least part of the wild-type gene encoding a

regulatory subunit of PI3K, and/or

(c) a second DNA sequence corresponding to at least part of the gene encoding a regulatory subunit of PI3K and containing a mutation of at least one nucleotide, which mutation corresponds to the mutation the presence of which in said gene is to be detected.

32. A test kit according to claim 31, wherein the second DNA sequence is a nucleic acid sequence according to any of claims 2-5.

10 33. A test kit according to claim 32, which further comprises means for amplifying DNA.

34. A test kit according to any of claims 31-33 for determining predisposition to impaired glucose tolerance in a subject.

15 35. A test kit according to the previous claim wherein the impaired glucose tolerance is related to an impaired glucose disappearance constant in a subject.

36. A test kit according to any of claims 31-33 for determining predisposition to decreased glucose effectiveness.

20

37. A test kit according to any one of claims 31-33 for determining predisposition to decreased insulin sensitivity in a subject.

38. A test kit for detecting the presence of a mutation in the gene encoding a regulatory
25 subunit of PI3K, the kit comprising

(a) means for amplifying DNA, and

(b) a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least
30 part of the gene encoding PI3K and containing a mutation of at least one nucleotide, which mutation corresponds to the mutation the presence of which in the gene encoding PI3K is to be detected.

39. A test kit according to claim 38, wherein the labelled oligonucleotide probe comprises a DNA isolate according to any of claim 2-5.

5 40. A test kit according to claim 38, which further comprises a labelled oligonucleotide probe comprising a DNA sequence corresponding to at least part of the wild-type gene encoding PI3K.

41. A test kit according to any of claims 38-40, wherein the label is selected from the group
10 consisting of enzymes, coloured or fluorescent substances, and radioactive isotopes.

42. Use of a nucleic acid sequence according to any one of claims 1-5 or a segment thereof as a probe, preferably an oligonucleotide probe, for detection of the presence of a mutation in a regulatory subunit of the enzyme PI3K.

15

43. Use according to the previous claim wherein the regulatory subunit is the human p85 α subunit or any isoform or homologous polypeptide thereof.

44. Use according to any one of claims 42-43, wherein the regulatory subunit is preferably
20 isolated from a biological sample obtained from a subject.

45. A variant of the p85 α regulatory subunit of human PI3K containing at least one amino acid substitution.

25 46. A variant according to the previous claim, in which methionine³²⁶ is substituted by isoleucine.

1 / 2




Fig. 1

SUBSTITUTE SHEET (RULE 26)

2/2




Fig. 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00200

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C12N 9/12, C12N 15/54, A01K 67/027, C12Q 1/48, C12Q 1/68, A61K 38/45
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, MEDLINE, BIOSIS, DBA, CA, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	Dialog Information Service, File 154, MEDLINE, Dialog accession no. 08999990, Medline accession no. 97184306, Hansen T et al: "Identification of a common amino acid polymorphism in the p85alpha regulatory subunit of phosphatidylinositol 3-kinase: effects on glucose disappearance constant, glucose effectiveness, and the insulin sensitivity index"; & Diabetes (UNITED STATES) Mar 1997, 46 (3) p494-501 --	1-46
P,X	Dialog Information Service, File 3, SciSearch, Dialog accession no. 14981552, Hansen T et al: "A missense mutation at codon 326(MET->ILE) in the gene encoding the phosphatidylinositol 3-kinase P85-alpha subunit is associated with decreased glucose-tolerance and decreased glucose effectiveness"; & Diabetologia, 1996, V39, S1 (AUG), P12 --	1-46

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

6 August 1997

Date of mailing of the international search report

14 -08- 1997

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Patrick Andersson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00200

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9612024 A1 (MAXPLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V.), 25 April 1996 (25.04.96), claims	1-2,4,6-10, 13-46
A	--	3,5,11-12
X	WO 9213001 A1 (NEW YORK UNIVERSITY), 6 August 1992 (06.08.92), figure 4, claim 19	1-2,4,6-10, 13-46
A	--	3,5,11-12
X	Proc. Natl. Acad. Sci., Volume 91, 1994, Kenta Hara et al, "1-Phosphatidylinositol 3-kinase activity is required for insulin stimulated glucose transport but not for RAS activation in CHO cells" page 7415 - page 7419	1-2,4,6-10, 13-46
A	-----	3,5,11-12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00200

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7-8 (partially)
because they relate to subject matter not required to be searched by this Authority, namely:

The search has been restricted to non-human living systems as the living system of the claims 7-8 may be a human, a subject which is generally accepted not to be patentable, see PCT search guidelines chapter VIII 1.1, line 6-8
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/07/97

International application No.

PCT/DK 97/00200

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9612024	A1	25/04/96	DE	4445562 C	04/04/96
<hr/>						
WO	9213001	A1	06/08/92	AU	667803 B	18/04/96
				AU	1234692 A	27/08/92
				CA	2100860 A	19/07/92
				EP	0567567 A	03/11/93
				JP	6505561 T	23/06/94
				US	5434064 A	18/07/95
				US	5618691 A	08/04/97
<hr/>						

THIS PAGE BLANK (USPTO)